

# MER 2003 Landing Sites– Impact Crater Lake and Hydrothermal Deposits

Horton Newsom,

Charles Barber, Ivan Thorsos


Institute of Meteoritics, Dept. of Earth and  
Planetary Sciences, University of New Mexico  
and Ashley Davies, Jet Propulsion Laboratory

# Method

- Determine craters that could contribute material to the landing sites.
- Obtain crater diameters using GIS.
- Determine hydrothermal potential for candidate craters.
- For superimposed craters calculate ejecta contributions to landing site ellipses.
- Determine fluvial contributions to landing site ellipses if applicable.



# Gusev Crater Site (MER-A)



# Hydrothermal deposits in Gusev

Gusev 158 km diam., melt - 10,000 km<sup>3</sup>


Thyra 21 km diam., melt - 140 km<sup>3</sup>

Gusev heat equivalent to 57 times  
Yellowstone over 15,000 years

- Ellipse contains Gusev rim material ejected from large superimposed craters including Thyra (but material is probably buried).
- Hydrothermal deposits are probably exposed on the rim of Thyra, inside of ellipse.
- Gusev rim material was probably transported to the ellipse by late fluvial processes.




MOLA  
Shaded  
Relief



# Transport of Material from Thyra to MER Landing Site


- Distance from center of landing site to Thyra's rim is only 21 km.
- Greater than 200 meter difference in elevation from the rim of Thyra down to the landing site allows for possible fluvial transport of material.

# Hematite site





**MOLA  
Shaded  
Relief**




Edge of  
hematite layer



# Ejecta transport

Viking  
MDIM




Superimposed 19  
km diam. crater  
H4 excavates  
from 2 km depth

MOC Wide Angle

Edge of ellipse

Edge of visible  
ejecta

Churning of  
surface by distal  
ejecta reveals  
hematite  
signature



# Hydrothermal deposits in Hematite site

H4 140 km diam., melt - 7,000 km<sup>3</sup>

Heat equivalent to 38 times

Yellowstone over 15,000 years

- Center of ellipse contains 20 to 40 cm of ejecta from superimposed 19 km diam. H5, including:
  - Impact melt from H4
  - Lake deposits in H4 (gravity -75 mgal, Yuan)
  - H4 rim material transported to H5 site by earlier superimposed craters.




# Isidis landing site: upstream craters

Courtesy L. Crumpler

## IS1 Crater

- 60 km diam.
- 500 km<sup>3</sup> impact melt
- 3 Yellowstone heat equivalents



# Conclusions – Lacustrine and Hydrothermal deposits

## ■ Lacustrine:

- Gusev (most likely) in surface layers
- Hematite site- in ejecta blanket of 19 km crater?
- Isidis - very unlikely

## ■ Hydrothermal:

- Hematite site (most likely) in ejecta blanket of 19 km crater
- Gusev – on rim of Thyra and in fluvial deposits
- Isidis – in fluvial deposits?





# TES Thermal Inertia on Geolines

